

# Journal Pre-proof

False-Negative COVID-19 Testing: Considerations in Obstetrical Care

Jeannie C. Kelly, MD MS, Michael Dombrowski, MD, Micaela O'neil-Callahan, MD, Annessa S. Kernberg, MD, Antonina I. Frolova, MD PhD, Molly J. Stout, MD MSCI



PII: S2589-9333(20)30073-2

DOI: <https://doi.org/10.1016/j.ajogmf.2020.100130>

Reference: AJOGMF 100130

To appear in: *American Journal of Obstetrics & Gynecology MFM*

Received Date: 17 April 2020

Accepted Date: 23 April 2020

Please cite this article as: Kelly JC, Dombrowski M, O'neil-Callahan M, Kernberg AS, Frolova AI, Stout MJ, False-Negative COVID-19 Testing: Considerations in Obstetrical Care, *American Journal of Obstetrics & Gynecology MFM* (2020), doi: <https://doi.org/10.1016/j.ajogmf.2020.100130>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Elsevier Inc. All rights reserved.

1                   **False-Negative COVID-19 Testing: Considerations in Obstetrical Care**

2           Jeannie C. KELLY MD MS<sup>1</sup>; Michael DOMBROWSKI MD<sup>1</sup>; Micaela O'NEIL-CALLAHAN MD<sup>2</sup>;

3           Annessa S. KERNBERG MD<sup>1</sup>; Antonina I. FROLOVA MD PhD<sup>1</sup>, Molly J. STOUT MD MSCI<sup>1</sup>

4           1. *Division of Maternal-Fetal Medicine, Washington University in St. Louis, St. Louis, Missouri*

5           2. *Affinia Healthcare, St. Louis, Missouri*

6  
7           The authors report no conflicts of interest.

8  
9           Correspondence should be addressed to: Jeannie C. Kelly, MD, MS. Washington University  
10          Department of Obstetrics and Gynecology, 660 S. Euclid Avenue, Campus Box 8064, St. Louis,  
11          MO 63110. Telephone: (314) 747-6788. Fax: 314 884-6007. Email: [jckelly@wustl.edu](mailto:jckelly@wustl.edu)

12  
13          **Word count:** Abstract 79 words; Manuscript 1,404 words

1 **Short title:** False-Negative COVID-19 testing

2

3 **Keywords:** COVID-19, SARS-CoV-2, pregnancy, coronavirus, diagnostic test sensitivity

Journal Pre-proof

4 **Introduction:** Real-time reverse transcriptions-polymerase chain reaction (RT-PCR) of  
5 nasopharyngeal (NP) swabs for SARS-Cov-2 are most commonly used for diagnosis of COVID-19  
6 infection, but there is limited information regarding the diagnostic test characteristics including  
7 negative and positive predictive values, including in pregnancy.

8

9 **Case:** A primiparous woman\* at 33 weeks' gestation presented to the obstetrical triage unit  
10 complaining of contractions, emesis, and cough for two days. She had fever, tachycardia,  
11 tachypnea, lymphopenia and mild elevation of liver enzymes. The fetus had reassuring testing,  
12 and her cervix was closed. Her BMI was 37.1 kg/m<sup>2</sup>, with no other co-morbidities. A chest  
13 radiograph showed subsegmental atelectasis without consolidation. Blood cultures, respiratory  
14 virus panel, and a NP swab for SARS-CoV2 PCR testing were sent. Empiric antibiotic therapy was  
15 initiated.

16

17 It was noted that her admission NP SARS-CoV2 PCR test obtained on day 3 of symptom was  
18 inadvertently sent out to a national reference laboratory, and thus a second test was  
19 performed on day 4 of symptom in-hospital for more timely results. Both tests resulted  
20 negative on that same day. Chest computed tomography revealed bilateral areas of  
21 consolidation and ground-glass opacification (Figure). All other infectious test results were  
22 negative. A third NP SARS-CoV2 PCR was obtained by the ICU staff on day 4 of symptoms, in  
23 case the prior two tests obtained by the obstetrical staff were limited by inadequate sampling.  
24 This test also resulted as negative the next day. The patient's cardiopulmonary status further  
25 worsened, and she was intubated. Given persistent maternal tachycardia at 150-160 bpm and

26 high fever requiring increasing amounts of vasopressor support, and fetal heart tracing with  
27 minimal variability, the team proceeded with a primary cesarean delivery. The neonate had  
28 Apgar scores of 1, 6, and 7 at 5, 10, and 15 minutes, respectively.

29 Bronchoalveolar lavage (BAL) performed after intubation by the ICU team revealed negative  
30 mycobacteriology and acid-fast stain, respiratory panel PCR, legionella culture, cytomegalovirus  
31 PCR, aerobic culture and gram stain, and adenovirus PCR; however, SARS-CoV2 RT-PCR of the  
32 BAL returned positive.

33 The patient remained intubated and in critical condition for 11 days. At the time of writing, she  
34 has been successfully extubated and transferred to a COVID-designated floor. The neonate is in  
35 good condition on room air in the neonatal ICU. NP SARS-CoV2 RT-PCR performed on the  
36 neonate on day of life 5 resulted negative.

37

38 **Discussion:** Three separate NP SARS-CoV2 RT-PCR tests from two institutions resulted as  
39 negative for a patient who was critically ill with a constellation of symptoms and lab findings  
40 consistent with COVID-19 infection, suggesting that false-negative testing is a clinically relevant  
41 problem not limited to a single platform with current testing strategies. In the non-pregnant  
42 population, sources of variability in RT-PCR testing results include the anatomic area sampled,  
43 quantity of virus present, stability of the RNA, timepoint in disease course, and assay  
44 variability.<sup>1-3</sup> False-negative result ranges of 17-63% for NP SARS-CoV2 RT-PCR have been  
45 reported in non-pregnant patients (Table); however, without clear gold standard tests available,  
46 diagnostic test characteristics including sensitivity, specificity, positive and negative predictive  
47 values of SARS-CoV2 RT-PCR assays are difficult to determine.<sup>1-3</sup> Sensitivity of BAL samples

48 appear to be higher than nasopharyngeal or oropharyngeal swabs, but requires invasive and  
49 high-risk aerosolizing bronchoscopy to obtain a sample.<sup>2,3</sup>

50

51 False-negative testing of NP SARS-CoV2 RT-PCR is a clinically relevant problem with multiple  
52 important implications, especially in pregnant women with suspicion for severe/critical COVID-  
53 19 infection: 1) Repeating NP SARS-CoV2 RT-PCR testing may be required for a positive result,  
54 as much as 3-5 times; 2) BAL SARS-CoV2 testing, a high-risk procedure, can be performed after  
55 negative NP SARS-CoV2 results if there is high clinical suspicion of COVID-19 infection and  
56 diagnosis is required for disposition; 3) Initially negative test results should not change clinical  
57 management; 4) Protocols should not allow for removal of precautions with a negative SARS-  
58 CoV2 test if there is high suspicion of COVID-19 infection; 5) All NP swab testing should  
59 performed by a specialized team, if possible, to improve uniformity in collection technique; 6) A  
60 universal testing strategy cannot be used as the single solution to risk stratify patients and  
61 determine infection prevention measures; 7) true population estimates of the disease are likely  
62 much underestimated,.

63 The most prudent strategy may be to presume that all patients are infected and use the best  
64 available infection prevention possible during the duration of this pandemic.

65

66

67 \*The patient's age was omitted to protect her identity.

68

69

70 **References**

- 71 1. Fang Y, Zhang H, Xie J, et al. Sensitivity of Chest CT for COVID-19: Comparison to RT-PCR.  
72 Radiology. February 2020:200432. doi:10.1148/radiol.2020200432
- 73 2. Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in Different Types of Clinical  
74 Specimens. JAMA. March 2020. doi:10.1001/jama.2020.3786
- 75 3. Yang Y, Yang M, Shen C, et al. Evaluating the Accuracy of Different Respiratory Specimens in  
76 the Laboratory Diagnosis and Monitoring the Viral Shedding of 2019-NCoV Infections.  
77 Infectious Diseases (except HIV/AIDS); 2020. doi:10.1101/2020.02.11.20021493
- 78 4. Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic proportion of  
79 coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship,  
80 Yokohama, Japan, 2020. Euro Surveill. 2020;25(10). doi:10.2807/1560-  
81 7917.ES.2020.25.10.2000180

82

83 **Figure 1.** Axial and coronal computed tomography images of the chest demonstrating severe  
84 bilateral disease.

86 **CRedit author statement**

87 **Kelly:** conceptualization, investigation, writing – original draft/editing/review & editing

88 **Dombrowski:** conceptualization, writing- review & editing, supervision **O’neil-Callahan:**

89 writing- review & editing, investigation **Kernberg:** software, writing- review & editing

90 investigation **Frolova:** writing- review & editing, investigation **Stout:** conceptualization,

91 investigation, writing– original draft/editing/review & editing, supervision

**Table: Current reports of false-negative RT-PCR nasal and/or pharyngeal swab testing for COVID-19**

Author	Country of origin	Study design	Primary aim	Total N	False negatives (%)	Positive on 1st test (%)	Positive on 2nd test (%)	Positive on 3rd test (%)	Maximum number of tests to obtain positive
Xiao <sup>1</sup>	China	Case series	Review of all RT-PCR tests that turned positive after initial negative test in one hospital	70	70 (100)	0 (0)	55 (78.6)	15 (21.4)	3
Ai <sup>2</sup>	China	Retrospective cohort	Comparison of chest CT with RT-PCR	1014	250* (24.7)	601 (59)	NS	NS	NS
Long <sup>3</sup>	China	Retrospective cohort	Comparison of chest CT with RT-PCR	36	6 (16.7)	30 (83.3)	3 (8.3)	3 (8.3)	3
Li <sup>4</sup>	China	Retrospective cohort	Review of RT-PCR tests in all patients diagnosed with COVID-19 by chest CT in one hospital	610	384 (63.0)	168 (27.5)	48 (7.9)	7 (1.1)	5
Wang <sup>5</sup>	China	Case report	Case report from Beijing	1	1 (100)	0 (0)	0 (0)	0 (0)	BAL required
Guo <sup>6</sup>	China	Retrospective cohort	Comparison of serum antibody testing with RT-PCR	208	58 (27.9)	NS	NS	NS	NS
Chen <sup>7</sup>	China	Case report	Case report from Hangzhou	1	1 (100)	0 (0)	1 (100)	0	2
Li <sup>8</sup>	China	Case series	2-patient case series from Beijing	2	2 (100)	0 (0)	1 (50%)	1 (50%)	2
Feng <sup>9</sup>	China	Case report	Case report from Zigong	1	1 (100)	0 (0)	0 (0)	0 (0)	5
Fang <sup>10</sup>	China	Retrospective cohort	Comparison of chest CT with RT-PCR	51	15 (29.4)	36 (70.6)	12 (23.5)	2 (3.9)	4
Wang <sup>11</sup>	China	Retrospective cohort	Comparison of RT-PCR results in different anatomical samples of confirmed cases	Nasal: 8 Pharyngeal: 398	Nasal: 3 (37.5) Pharyngeal: 272 (68.3)	NS	NS	NS	NS
Yang <sup>12</sup>	China	Retrospective cohort	Comparison of RT-PCR results in different anatomical samples and time points of confirmed cases**	Nasal: 445 Throat: 158	Nasal: 157 (35.3) Throat: 74 (46.8)	NS	NS	NS	NS

NS: not specified

BAL: bronchoalveolar lavage

Author names italicized for publications also referenced in manuscript (10-12)

\*Based on CT-scan findings and clinical correlation

\*\* Results from 14 days of symptom onset included

1. Xiao AT, Tong YX, Zhang S. False-negative of RT-PCR and prolonged nucleic acid conversion in COVID-19: Rather than recurrence. *Journal of Medical Virology*. doi:10.1002/jmv.25855
2. Ai T, Yang Z, Hou H, et al. Correlation of Chest CT and RT-PCR Testing in Coronavirus Disease 2019 (COVID-19) in China: A Report of 1014 Cases. *Radiology*. February 2020:200642. doi:10.1148/radiol.2020200642
3. Long C, Xu H, Shen Q, et al. Diagnosis of the Coronavirus disease (COVID-19): rRT-PCR or CT? *Eur J Radiol*. 2020;126:108961. doi:10.1016/j.ejrad.2020.108961



4. Li Y, Yao L, Li J, et al. Stability issues of RT-PCR testing of SARS-CoV-2 for hospitalized patients clinically diagnosed with COVID-19. *Journal of Medical Virology*. n/a(n/a).  
doi:10.1002/jmv.25786
5. Wang Y, Kang H, Liu X, Tong Z. Combination of RT-qPCR testing and clinical features for diagnosis of COVID-19 facilitates management of SARS-CoV-2 outbreak. *Journal of Medical Virology*. 2020;92(6):538-539. doi:10.1002/jmv.25721
6. Guo L, Ren L, Yang S, et al. Profiling Early Humoral Response to Diagnose Novel Coronavirus Disease (COVID-19). *Clin Infect Dis*. doi:10.1093/cid/ciaa310
7. Chen Z, Li Y, Wu B, Hou Y, Bao J, Deng X. A Patient with COVID-19 Presenting a False-Negative Reverse Transcriptase Polymerase Chain Reaction Result. *Korean Journal of Radiology*. 2020;21. doi:10.3348/kjr.2020.0195
8. Li D, Wang D, Dong J, et al. False-Negative Results of Real-Time Reverse-Transcriptase Polymerase Chain Reaction for Severe Acute Respiratory Syndrome Coronavirus 2: Role of Deep-Learning-Based CT Diagnosis and Insights from Two Cases. *Korean Journal of Radiology*. 2020;21(4):505-508. doi:10.3348/kjr.2020.0146
9. Feng H, Liu Y, Lv M, Zhong J. A case report of COVID-19 with false negative RT-PCR test: necessity of chest CT. *Jpn J Radiol*. April 2020. doi:10.1007/s11604-020-00967-9
10. Fang Y, Zhang H, Xie J, et al. Sensitivity of Chest CT for COVID-19: Comparison to RT-PCR. *Radiology*. February 2020:200432. doi:10.1148/radiol.2020200432
11. Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in Different Types of Clinical Specimens. *JAMA*. March 2020. doi:10.1001/jama.2020.3786

12. Yang Y, Yang M, Shen C, et al. Evaluating the Accuracy of Different Respiratory Specimens in the Laboratory Diagnosis and Monitoring the Viral Shedding of 2019-NCoV Infections. *Infectious Diseases (except HIV/AIDS)*; 2020. doi:10.1101/2020.02.11.20021493

Journal Pre-proof

